

SHORTER CONTRIBUTIONS

Banisteria, Number 27, 2006

© 2006 by the Virginia Natural History Society

HEMIEPLUS MICROPHTHALMUS (SCHWARZ): A SPECIES AND GENUS OF ELONGATED BEETLES NEW TO THE FAUNA OF VIRGINIA (COLEOPTERA: MYCTERIDAE) – Ongoing faunistic inventories conducted in southeastern Virginia continue to disclose a wide variety of insects substantially north of their previously-known geographic ranges. The latest addition to this category of overlooked native Virginians is *Hemipeplus microphthalmus*, a small, flattened beetle with remarkably elongated abdomen, described by E. A. Schwarz (1878) from Enterprise, Florida, and subsequently found frequently in Florida and as single collections in Mississippi, Texas, and South Carolina (Thomas, 1985).

Discovery of this species in Virginia extends the known range along the coast some 300 miles/485 km north from Georgetown, South Carolina, and represents a new northernmost locality for this basically tropical genus, two other species of which occur in Florida and southern Georgia. Collection data for the specimen are:

Virginia: *City of Virginia Beach:* False Cape State Park, Wash Woods Trail, 0.3 km east of main park road, UV trap in live oak–loblolly pine dune woods, 2 August 2005, S. M. Roble and G. W. Wahl, III (VMNH).

Referring to the habits of this species in Florida, Thomas (1985) stated: “In contrast to the other two species of Florida *Hemipeplus*, this species does not seem to be associated with palmetto. Instead, it occurs on grasses of various kinds, dwelling between the closely appressed leaf bases and stem.” One host plant mentioned by Thomas was *Andropogon virginicus* L. (Poaceae) which is statewide in Virginia (Harvill et al., 1992). Attention directed specifically to this plant at False Cape should reveal the extent to which it is utilized by *H. microphthalmus* there, and elsewhere in the coastal regions of the state. Interestingly, *H. microphthalmus* is uniformly light yellow-tan and doubtless matches the color of *Andropogon* foliage closely. There is no evidence that the beetles actually feed on the plant tissue, and Thomas believed that they are scavengers on fungi and miscellaneous plant spores.

An interesting parallel modification of body form occurs in many laminaphilic species of the lygaeid subfamily Blissinae. For example, *Ischnodesmus falicus* (Say), a species that occurs in Virginia, is very flat and slender with the abdomen greatly elongated: 58% of the total body length (Hoffman, 1996: fig. 29). Insects adapted to life in the axils of grass blades are not




Fig. 1. *Hemipeplus microphthalmus*, dorsal aspect of specimen from False Cape State Park. Scale line = 1 mm.

frequently collected by conventional sweeping. Perhaps a quantity of the plants processed with a Berlese extraction funnel might yield more material of *H. microphthalmus*, as the species is not especially phototropic.

Because of the singular body form, *Hemipeplus* was until fairly recently (e.g., Arnett, 1963) placed in the family Cucujidae, a family with many flattened subcorticolous species. During that time it sometimes formed the basis of its own family, or was associated with a variety of beetle families. Thomas' (1985) review recounted in detail the taxonomic and nomenclatorial convolutions through which the genus has passed. Its current placement in the Mycteridae is predicated largely

upon larval characters: adult beetles of *Hemipeplus* and of *Mycterus* reflect few features in common beyond those of the entire order.

ACKNOWLEDGMENTS

The Virginia Museum of Natural History is obliged to Dr. Steven M. Roble (Division of Natural Heritage, Virginia Department of Conservation and Recreation) for transfer of collections resulting from its inventory activities, including that in which the specimen of *Hemipeplus* was found. I express also my gratitude to Dr. Michael C. Thomas (Florida State Collection of Arthropods) for identifying the specimen and suggesting useful literature resources.

LITERATURE CITED

Arnett, R. H. Jr. 1960-1963. The Beetles of the United States (A Manual for Identification). Catholic University of America Press, Washington. 1,112 pp.

Harvill, A. M., T. R. Bradley, C. E. Stevens, T. F. Wieboldt, D. M. E. Ware, D. W. Ogle, G. W. Ramsey, & G. P. Fleming. 1992. Atlas of the Virginia Flora. Virginia Botanical Associates, Burkeville, VA. 144 pp.

Hoffman, R. L. 1996. Seed bugs of Virginia (Heteroptera: Lygaeidae). *Insects of Virginia* 14: 1-111.

Schwarz, E. A. 1878. The Coleoptera of Florida. *Proceedings of the American Philosophical Society* 17: 353-469.

Thomas, M. C. 1985. The species of *Hemipeplus* Latreille (Coleoptera: Mycteridae) in Florida, with a taxonomic history of the genus. *The Coleopterists Bulletin* 39: 365-375.

Richard L. Hoffman
Virginia Museum of Natural History
Martinsville, Virginia 24112

Banisteria, Number 27, 2006
© 2006 by the Virginia Natural History Society

A NOTE ON THE OCCURRENCE OF THE SPIDER *ARACHOSIA CUBANA* (BANKS) IN VIRGINIA (ARANEAE: ANYPHAENIDAE) – My recent synopsis (Hoffman, 2002) of the Virginia species of anyphaenid spiders treated this species under the name *Oxysoma*

cubana, provided two new locality records for the Coastal Plain, and cast doubt on an earlier published record by Howell & Pienkowski (1971) for Montgomery County. The notably scarce material available to Platnick (1974) when he revised the family implied a basically maritime distribution in eastern United States. Montgomery County, Virginia thus seemed to me “biogeographically improbable.” Recently acquired specimens of this species from two localities in southwestern Virginia are noteworthy in confirming the presence of *A. cubana* in the mountains of the state and cancelling my unjustified suspicion of the previous record. To the known localities, I can now add:

Floyd County: sedge meadow beside US 221, 2 km west of Willis, 17-18 June 2004, S. M. Roble (VMNH, 1 m), and *Grayson County*: New River at the US 221/58 bridge, southwest of Galax, 9 June 2005, S. M. Roble (VMNH, 1m). These specimens differ slightly from Coastal Plain males in lacking a black stripe at the base of each leg spine, but agree closely in details of male genitalia.

Although *A. cubana* is easily recognizable as an anyphaenid (general habitus, clustered eyes, lamellate paronychia hair tufts), it differs strikingly from the other Nearctic genera of the family in lacking any trace of the retrolateral tibial apophysis of the male pedipalp. Ramirez (2003) transferred the species from *Oxysoma* to *Arachosia*, as the only local member of the subfamily Amaurobioidinae.

LITERATURE CITED

Howell, J. O., & R. L. Pienkowski. 1971. Spider populations in alfalfa, with notes on spider prey and effect of harvest. *Journal of Economic Entomology* 64: 163-168.

Hoffman, R. L. 2002. Spiders of the family Anyphaenidae in Virginia. *Banisteria* 19: 9-12.

Platnick, N. I. 1974. The spider family Anyphaenidae in America north of Mexico. *Bulletin of the Museum of Comparative Zoology* 146: 205-266.

Ramirez, M. J. 2003. The spider subfamily Amaurobioidinae (Araneae, Anyphaenidae): a phylogenetic revision at the generic level. *Bulletin of the American Museum of Natural History* 277: 1-262.

Richard L. Hoffman
Virginia Museum of Natural History
Martinsville, Virginia 24112